arXiv:1401.7151v2 [math.OA] 27 Apr 2014

CONTINUOUS FIELDS OF POSTLIMINAL C^*-ALGEBRAS

ALDO J. LAZAR

Abstract. We discuss a problem of Dixmier [6, Problem 10.10.11] on continuous fields of postliminal C^*-algebras and the greatest liminal ideals of the fibers.

1. Introduction

In [6, Problem 10.10.11] Dixmier asked the following question: given a continuous field $((A(t)), \Theta)$ of postliminal C^*-algebras over some topological space T and with $B(t)$ the greatest liminal ideal of $A(t)$ and $\Theta' := \{x \in \Theta \mid x(t) \in B(t), t \in T\}$ is $((B(t)), \Theta')$ a continuous field of C^*-algebras? We shall call a continuous field of postliminal -âlgebras for which the answer to this question is affirmative a tame continuous field.

An example of a continuous field that is not tame can be constructed over $T := \mathbb{N} \cup \{\infty\}$. We let $A(n)$, $n \in \mathbb{N}$, be the unitization of $K(H)$, the algebra of all compact operators over an infinite dimensional Hilbert space H, and $A(\infty) := \mathbb{C}I_H$, I_H being the identity operator on H. Θ consists of all the fields x such that $x(n) = \lambda_n I_H + a_n$, $\{\lambda_n\}$ being a sequence in $\mathbb{C}$ that converges to some $\lambda \in \mathbb{C}$, $\{a_n\}$ being a sequence in $K(H)$ that converges to $\{0\}$, and $x(\infty) = \lambda I_H$. Then $((A(t))_{t\in T}, \Theta)$ is a continuous field of postliminal C^*-algebras. Now the largest liminal ideal of $A(n)$ is $B(n) = K(H)$ and the largest liminal ideal of $A(\infty)$ is $B(\infty) = A(\infty) = \mathbb{C}I_H$. Clearly $x \in \Theta$ satisfies $x(t) \in B(t)$ for every $t \in T$ only if $x(\infty) = 0$ and the continuous field is not tame.

In the next section we shall show that the continuous fields of postliminal C^*-algebras in a certain class that properly includes the locally trivial continuous fields

Date: April 22, 2014.

1991 *Mathematics Subject Classification.* 46L05.

Key words and phrases. continuous fields of C^*-algebras, postliminal C^*-algebras, liminal C^* algebras.

We wish to express our gratitude to Professor E. Christensen and Professor J. Phillips for providing us the reference [4].

are always tame. Another result which we prove is a necessary and sufficient condition for tameness expressed in terms related to the minimal primitive ideals of the fibers. Afterwards we shall exhibit an example of a continuous field of postliminal C^*-algebras such that all its fibers are mutually isomorphic and its restriction to any open subset of the base space is not tame.

Let A be the C^*-algebra of the continuous field $((A(t)), \Theta)$ of C^*-algebras over the locally compact Hausdorff space T as defined in [6, 10.4.1]. By [7, Theorem 1.1] for every primitive ideal P of A there exist a unique $t_P \in T$ and a unique primitive ideal Q_P of $A(t_P)$ such that $P = \{x \in A \mid x(t_P) \in Q_P\}$ and conversely every pair (t, Q), $t \in T$, $Q \in \mathrm{Prim}(A(t)$ determines a primitive ideal of A in this manner. Obviously P is minimal in $\mathrm{Prim}(A)$ if and only if Q_P is minimal in $A(t_P)$.

We shall use the terminology and the notation for continuous fields as introduced in [6, Chapter 10]. The preference to work with continuous fields rather than Banach bundles as it is more common nowadays is motivated by the fact that Dixmier's original question was expressed in these terms. The closed unit ball of the Banach space X is denoted X_1.

2. Results

The main ingredient in the proof of Proposition 1 that follows is Michael's selection theorem [10, Theorem 3.2"]: a multivalued map φ from a paracompact space T to the family of the non-void closed convex subsets of a Banach space X that is lower semicontinuous admits a continuous selection, i.e., there is a continuous function $f : T \to X$ such that $f(t) \in \varphi(t)$ for every $t \in T$. Moreover, if F is a closed subset of T and $g : F \to X$ is a continuous selection for $\varphi \mid_F$ then one may choose f so $f \mid_F = g$ is satisfied. Recall that φ is called lower semicontinuous if for each open subset U of X the set $\{t \in T \mid \varphi(t) \cap U \neq \emptyset\}$ is open.

Proposition 1. *Let T be a paracompact space or a locally compact Hausdorff space and X a Banach space. Denote by $\mathcal{M}$ the space of the closed unit balls of all the closed subspaces of X endowed with the Hausdorff metric. Suppose $t \to X(t)_1$, $t \in T$, is a continuous map into $\mathcal{M}$, $X(t)$ being a closed subspace of X. With Γ the space of all the continuous functions $\varphi : T \to X$ such that $\varphi(t) \in X(t)$, $t \in T$, $((X(t)), \Gamma)$ is a continuous field of Banach spaces.*

Proof. The only evidence we must provide is that for $t_0 \in T$ and $x_0 \in X(t_0)$ there exists $\varphi \in \Gamma$ such that $\varphi(t_0) = x_0$. Clearly we may suppose $x_0 \neq 0$. Set $y_0 := x_0/\|x_0\|$. We claim that $t \to X(t)_1$ is lower semicontinuous as a multivalued map from T to X. To see this let U be an open subset of X, $s \in \{t \in T \mid U \cap X(t)_1 \neq \emptyset\}$, and $z \in U \cap X(s)_1$. There are an open ball of X of center z and of radius $\varepsilon > 0$ contained in U and a neighborhood V of s in T such that $d(X(s)_1, X(t)_1) < \varepsilon$ for all $t \in V$, d being the Hausdorff metric. Thus for each $t \in V$ there is $w_t \in X(t)_1$ for which $\|z - w_t\| < \varepsilon$. It follows that $V \subset \{t \in T \mid U \cap X(t) \neq \emptyset\}$ and we conclude that $\{t \in T \mid U \cap X(t) \neq \emptyset\}$ is open. We obtained that the map $t \to X(t)_1$ is indeed lower semicontinuous.

Suppose now that T is paracompact. By Michael's selection theorem mentioned above there exists a continuous map $\varphi' : T \to X$ such that $\varphi'(t) \in X(t)$ for every $t \in T$ and $\varphi'(t_0) = y_0$. The map φ defined by $\varphi(t) := \|x_0\|\varphi'(t)$ suits the requirements.

Let now T be locally compact Hausdorff. Let W be a compact neighborhood of t_0. Again by Michael's selection theorem there is a continuous map $\varphi' : W \to X$ such that $\varphi'(t) \in X(t)$ for every $t \in W$ and $\varphi'(t_0) = y_0$. Let now $f : T \to [0, 1]$ be a continuous function such that $f(t_0) = 1$ and $f(t) = 0$ for $t \notin Int(W)$. The function $\varphi : T \to X$ defined by

$$\varphi(t) := \begin{cases} \|x_0\| f(t)\varphi'(t), & \text{if } t \in W, \\ 0, & \text{if } t \notin W. \end{cases}$$

is continuous, satisfies $\varphi(t) \in X(t)$ for $t \in T$ and $\varphi(t_0) = x_0$.

□

A continuous field of Banach spaces over a paracompact or a locally compact Hausdorff space T isomorphic to a continuous field of Banach spaces as described in Proposition 1 will be called uniform. Obviously, a trivial continuous field of Banach spaces is uniform. A continuous field of Banach spaces over T is called locally uniform if T has an open cover $\{U_\alpha\}$ such that its restriction to each U_α is uniform. By using the regularity of the base space as was done in the last paragraph of the previous proof one gets the following Proposition.

Proposition 2. *Let T be as in Proposition 1, $\{U_\alpha\}_{\alpha \in \mathcal{A}}$ an open cover of T and X_α a Banach space, $\alpha \in \mathcal{A}$. Denote by M_α the space of the closed unit balls of all the*

closed subspaces of X_α endowed with the Hausdorff metric. Suppose $X(t)$, $t \in T$, is a Banach space that is a closed subspace of X_α whenever $t \in U_\alpha$. Moreover, suppose that the map $t \to X(t)_1$ from U_α into M_α, $\alpha \in \mathcal{A}$, is continuous. Denote by Γ the space of all the functions $\varphi : T \to \cup_{\alpha\in\mathcal{A}} X_\alpha$ such that $\varphi(t) \in X(t)$, $t \in T$, and restriction of φ to U_α is continuous as a map into X_α, $\alpha \in \mathcal{A}$. Then $((X(t)), \Gamma)$ is a locally uniform continuous field of Banach spaces.

Of interest for us are the continuous fields of C^*-algebras; of course, for uniform continuous fields of C^*-algebras we shall require that the Banach space appearing in the definition be a C^*-algebra and the fibers to be C^*-algebras of it. It is natural to ask if a uniform continuous field of C^*-algebras has to be locally trivial. There is some indication in [4, Theorem 4.3] that this may be the case when the fibers are nuclear and separable. On the other hand, [3, Theorem 3.3] provides an example of a uniform continuous field of (non-separable) nuclear C^*-algebras that is not locally trivial.

Theorem 3. *Suppose $((A(t)), \Theta)$ is a locally uniform continuous field of postliminal C^*-algebras over a paracompact or locally compact Hausdorff space T. Let $B(t)$ be the largest liminal ideal of $A(t)$ and $\Theta' := \{x \in \Theta \mid x(t) \in B(t), t \in T\}$. Then $((B(t)), \Theta')$ is a locally uniform continuous field of C^*-algebras over T.*

Proof. Let U be an open subset of T over which $((A(t)), \Theta)$ is uniform. There is no loss of generality if we suppose that over U all the fibers $A(t)$ are C^*-subalgebras of a certain C^*-algebra and $t \to A(t)_1$ is continuous for the Hausdorff metric. If $t', t'' \in U$ satisfy $d(A(t')_1, A(t'')_1) < s(\leq 1/21)$ then it follows from [13, Lemma 1.10] and [12, Theorem 2.7] that $d(B(t')_1, B(t'')_1) < 7s$. Hence $t \to B(t)_1$ is continuous on U and the conclusion follows from Proposition 2.

□

We shall discuss now the behaviour of locally uniform continuous fields of postliminal C^*-algebras with respect to two kinds of ideals that give rise to canonical composition series. Recall that one says that a point π in the spectrum $\hat{A}$ of a C^*-algebra A satisfies the Fell condition if there exist a neighbourhood V of π in $\hat{A}$ and $a \in A^+$ such that $\varrho(a)$ is a projection of rank 1 for every $\varrho \in V$. A Fell C^*-algebra is a C^*-algebra for which all the points in its spectrum satisfy the Fell condition,

see [1] and [11, 6.1] where these algebras were called of Type I_0. Every non trivial postliminal C^*-algebra has a non trivial largest Fell ideal by [11, Proposition 6.1.7]. A C^*-algebra A is called uniformly liminal if its ideal of all the elements $a \in A$ for which the function $\pi \to \pi(a)$ is bounded on $\hat{A}$ is dense in A, see [2, p. 443] and the references given there. Every non trivial postliminal C^*-algebra has a non trivial largest uniformly liminal ideal by [2, Theorem 2.6 and Theorem 2.8].

Theorem 4. *Let $((A(t)), \Theta)$ be a locally uniform continuous field of postliminal C^*-algebras over a paracompact or locally compact Hausdorff space T. Let $B(t)$ be the largest Fell ideal of $A(t)$ and $\Theta' := \{x \in \Theta \mid x(t) \in B(t), t \in T\}$. Then $((B(t)), \Theta')$ is a locally uniform continuous field of C^*-algebras.*

Proof. Let U be an open subset of T such that the restriction of $((A(t)), \Theta)$ to U is uniform; to simplify matters we shall suppose that all the C^*-algebras $A(t)$, $t \in U$ are C^*subalgebras of a certain C^*algebra and the map $t \to A(t)_1$ is continuous on U for the Hausdorff metric. Let now $t_1, t_2 \in U$ satisfy $d(A(t_1)_1, A(t_2)_1) < s(< 1/147)$. By [13, Lemma 1.10] there is an ideal J of $A(t_2)$ such that $d(B(t_1)_1, J_1) < 7s$. The same lemma yields a homeomorphism h from $\hat{J}$ onto $\widehat{B(t_1)}$. Pick $\pi_0 \in \hat{J}$ and set $\varrho_0 = h(\pi_0) \in \widehat{B(t_1)}$. There is $a \in B(t_1)_1^+$ and a neighbourhood V of ϱ_0 in $\widehat{B(t_1)}$ such that $\varrho(a)$ is a rank 1 projection for $\varrho \in V$. The proof of [13, Lemma 2.4] yields an element $b \in B(t_2)_1^+$ such that $\pi(b)$ is a rank 1 projection when $\pi \in h^{-1}(V)$. It follows that J is a Fell ideal contained in $B(t_2)$. Similarly, there is a Fell ideal L of $A(t_1)$ such that $d(L_1, B(t_2)_1) < 7s$ which has to satisfy $B(t_1) \subset L$ by [13, Lemma 1.10]. Thus $L = B(t_1)$ and $d(B(t_1)_1 B(t_2)_1) < 7s$. We found that $t \to B(t)_1$ is continuous on U and Proposition 2 provides the conclusion. □

Very likely the hypothesis of separability in the next result is unnecessary but we were not able to find a proof that dispenses with it.

Theorem 5. *Let $((A(t)), \Theta)$ be a locally uniform continuous field of separable postliminal C^*-algebras over a paracompact or locally compact Hausdorff space T. Let $B(t)$ be the largest uniformly liminal ideal of $A(t)$ and $\Theta' := \{x \in \Theta \mid x(t) \in B(t), t \in T\}$. Then $((B(t)), \Theta')$ is a locally uniform continuous field of C^*-algebras.*

Proof. As in the previous proof, we consider an open subset U of T such that the given continuous field of C^*-algebras restricted to U is uniform. We shall also suppose that the family of C^*-algebras $\{A(t) \mid t \in U\}$ is contained in a certain C^*-algebra and the map $t \to A(t)_1$ is continuous on U for the Hausdorff metric. If $t_1, t_2 \in U$ satisfy $d(A(t_1)_1, A(t_2)_1) < s < 1/420000$ then $A(t_1)$ and $A(t_2)$ are isomorphic by [4, Theorem 4.3]. Of course, every isomorphism between these two C^*-algebras maps $B(t_1)$ onto $B(t_2)$. Hence, the same result of [4] tells us that $d(B(t_1), B(t_2)) < 28s^{1/2}$ so the map $t \to B(t)_1$ is continuous on U. Once more Proposition 2 yields the conclusion.

□

Question 6. *Let $((A(t)), \Theta)$ be a uniform continuous field of postliminal C^*-algebras over T. Can one choose a non-trivial continuous trace ideal $B(t)$, $t \in T$, of $A(t)$ such that with $\Theta' := \{x \in \Theta \mid x(t) \in B(t), t \in T\}$, $((B(t)), \Theta')$ is a continuous field of C^*-algebras?*

Given a continuous field $((A(t)), \Theta)$ of postliminal C^*-algebras over a locally compact Hausdorff space T, let A be the C^*-algebra defined by this continuous field; obviously it is a postliminal C^*-algebra. Let B be its greatest liminal ideal. If the image of B in $A(t)$ by the evaluation map is the greatest liminal ideal of $A(t)$, $t \in T$, then it is easily seen that the given field is tame. Conversely, suppose that $((A(t)), \Theta)$ is tame and let C be the C^*-algebra defined by the continuous field of the greatest liminal ideals. Then $C = B$ the greatest liminal ideal of A. Indeed, it is clear that C is a liminal ideal of A so $C \subset B$. Let now $x \in B$. With $t \in T$, $\rho \in \widehat{A(t)}$, we have that $y \to \rho(y(t))$, $y \in A$, is an irreducible representation of A hence $\rho(x(t))$ is a compact operator over the space of the representation. We conclude by [6, 4.2.6] that $x(t) \in C(t)$. Thus $B \subset C$.

Very likely the following lemma is known but in the absence of a reference for it we give its simple proof.

Lemma 7. *Let A be a postliminal C^*-algebra, $M \subset \mathrm{Prim}(A)$ the set of all the minimal primitive ideals. Then $\mathrm{Int}(M)$ is the primitive ideal space of the greatest liminal ideal I of A.*

Proof. Dixmier remarked in [5, Remarque C, p. 111] that we have $\mathrm{Prim}(I) \subset \mathrm{Int}(M)$. $\mathrm{Int}(M)$ is the primitive ideal space of an ideal, J say, of A. If $P \in \mathrm{Int}(M)$ then the relative closure of $\{P\}$ in $\mathrm{Int}(M)$ is just $\{P\}$ since all the ideals in $\mathrm{Int}(M)$ are minimal primitive ideals. Thus $\mathrm{Int}(M)$ is T_1 in the relative topology and we gather that J is a liminal ideal of A. Hence $J \subset I$ and $\mathrm{Int}(M) \subset \mathrm{Prim}(I)$

□

The set M of above need not be open; for examples see [8] and [9, Example 4.3].

We can now state and prove a necessary and sufficient condition for the tameness of a continuous field of postliminal C^*-algebras.

Theorem 8. *Let $\mathcal{C} := ((A(t)), \Theta)$ be a continuous field of postliminal C^*-algebras over the locally compact Hausdorff space T and A the C^*-algebra defined by $\mathcal{C}$. Denote by W the set of non minimal ideals in* $\mathrm{Prim}(A)$*, $W(t) := W \cap \mathrm{Prim}((A(t))$, $I(t) := \cap\{P \mid P \in W(t)\}$ for $t \in T$, and $\tilde{\Theta} := \{x \in \Theta \mid x(t) \in I(t), t \in T\}$. Then $\mathcal{C}$ is tame if and only if $((I(t)), \tilde{\Theta})$ is a continuous field of C^*-algebras.*

Proof. By Lemma 7 and the remarks preceding it $\mathcal{C}$ is tame if and only if $\mathrm{Prim}(A(t)) \cap \overline{W} = \overline{W(t)}$ for every $t \in T$. Let $s \in T$ and set $J(s) := \{x(s) \mid x \in \Theta, x(t) \in I(t)$ for all $t \in T\} \subset I(s)$. Then it follows from [7, Theorem 1.2] that $\mathrm{Prim}(A(s)) \cap \overline{W} = \mathrm{hull}(J(s))$. Now $\overline{W(t)} = \mathrm{hull}(I(t))$ so $\mathcal{C}$ is tame if and only if $\mathrm{hull}(J(t) = \mathrm{hull}(I(t))$ for each $t \in T$ if and only if $\overline{J(t)} = I(t)$ for each $t \in T$. This means that $\mathcal{C}$ is tame if and only if $((I(t)), \tilde{\Theta})$ is a continuous field of C^*-algebras and we are done.

□

3. An Example

As mentioned in the Introduction, we are going to construct in this section a continuous field of postliminal C^*-algebras over $[0, 1]$ whose fibers are mutually isomorphic and which has the additional property that none of its restrictions to the relatively open subsets of $[0, 1]$ is tame.

First we proceed to prepare two C^*-algebras that will serve as building blocks of the fibers. Let $\mathbb{N} = \cup_{p=1}^{\infty} S_p$ where the sets $\{S_p\}$ are mutually disjoint and each $S_p = \{n_1^p < n_2^p < \ldots\}$ is infinite. Let H be a separable Hilbert space with an

orthonormal basis $\{\xi_k\}_{k=1}^{\infty}$ and $\mathcal{B}(H)$ the C^*-algebra of all the bounded operators on H. Denote by e_{ij}^0 the partial isometry that maps ξ_j to ξ_i and vanishes on each ξ_k with $k \neq j$. The C^*-subalgebra of $\mathcal{B}(H)$ generated by $\{e_{ij}^0 \mid i,j = 1,2,\ldots\}$ is the ideal of all compact operators and we shall denote it by A_0. Put now $e_{ij}^1 := \sum_{m=1}^{\infty} e^0_{n_m^i n_m^j}$, $i,j = 1,2,\ldots$ where the series converges in the strong operator topology. Then $\{e_{ii}^1\}_{i=1}^{\infty}$ are mutually orthogonal projections, $\sum_{i=1}^{\infty} e_{ii}^1 = \mathbf{1}_H$, e_{ij}^1 is a partial isometry from $e_{jj}^1(H)$ onto $e_{ii}^1(H)$, $(e_{ij}^1)^* = e_{ji}^1$ and

$$e_{ij}^1 e_{rs}^1 = \begin{cases} e_{is}^1, & j = r \\ 0, & j \neq r. \end{cases}$$

Hence the C^*-subalgebra K_1 of $\mathcal{B}(H)$generated by $\{e_{ij}^1 \mid i,j = 1,2,\ldots\}$ is isomorphic to A_0 and $A_0 \cap K_1 = \{0\}$. We have

$$e_{ij}^1 e_{rs}^0 = \begin{cases} e^0_{n_m^i s}, & \text{if } r = n_m^j \text{ for some } m, \\ 0, & \text{otherwise} \end{cases} \tag{1}$$

and

$$e_{rs}^0 e_{ij}^1 = \begin{cases} e^0_{r n_m^j}, & \text{if } s = n_m^i \text{ for some } m, \\ 0, & \text{otherwise.} \end{cases} \tag{2}$$

$A_1 := A_0 + K_1$ is a postliminal C^*-algebra since A_0 and $K_1 \sim A_1/A_0$ are postliminal C^*-algebras (actually liminal C^*-algebras in this case). Each $x \in A_1$ admits a unique decomposition $x = x_0 + x_{K_1}$ with $x_0 \in A_0$ and $x_{K_1} \in K_1$. The map $x \to x_{K_1}$ is a homomorphism hence $\|x_{K_1}\| \leq \|x\|$ and $\|x_0\| \leq 2\|x\|$.

From (1) and (2) it follows that the sequence $\{\sum_{i=1}^{m} e_{ii}^1\}_{m=1}^{\infty}$ is an increasing approximate unit for A_1 consisting of projections.

We suppose now that the C^*-subalgebras $K_1, \ldots K_{l-1}$ of $\mathcal{B}(H)$ have been defined such that K_p, $1 \leq p \leq l-1$, is spanned by

$$e_{ij}^p := \sum_{m=1}^{\infty} e^{p-1}_{n_m^i n_m^j}, i,j = 1,2,\ldots.$$

It follows that $\{e_{ii}^p\}_{i=1}^{\infty}$ are mutually orthogonal projections, $\sum_{i=1}^{\infty} e_{ii}^p = \mathbf{1}_H$, and e_{ij}^p is a partial isometry from $e_{jj}^p(H)$ onto $e_{ii}^p(H)$. With $A_p := A_{p-1} + K_p$, $1 \leq p \leq l-1$, we have $A_{p-1} \cap K_p = \{0\}$, A_p is a postliminal C^*-subalgebra of $\mathcal{B}(H)$, A_{p-1} is an ideal of A_p and $\{\sum_{i=1}^{m} e_{ii}^p\}_{m=1}^{\infty}$ is an increasing approximate unit of A_p consisting of projections. Every element $x \in A_p$ admits a unique decomposition $x = x_{p-1} + x_{K_p}$ where $x_{p-1} \in A_{p-1}$, $x_{K_p} \in K_p$. Moreover, $x \to x_{K_p}$ is a homomorphism hence $\|x_{K_p}\| \leq \|x\|$ and $\|x_{p-1}\| \leq 2\|x\|$.

We define now

$$e^l_{ij} := \sum_{m=1}^{\infty} e^{l-1}_{n^i_m n^j_m}, i, j = 1, 2, \ldots.$$

Then $\{e^l_{ii}\}_{i=1}^{\infty}$ are mutually orthogonal projections, $\sum_{i=1}^{\infty} e^l_{ii} = \mathbf{1}_H$, and e^l_{ij} is a partial isometry from $e^l_{jj}(H)$ onto $e^l_{ii}(H)$. We obtain $(e^l_{ij})^* = e^l_{ji}$ and

$$e^l_{ij} e^l_{rs} = \begin{cases} e^l_{is}, & \text{if } j = r, \\ 0, & \text{otherwise,} \end{cases} \tag{3}$$

hence the C^*-subalgebra K_l of $\mathcal{B}(H)$ generated by $\{e^l_{ij} \mid i, j = 1, 2, \ldots\}$ is isomorphic to A_0. We also have

$$e^l_{ij} e^{l-1}_{rs} = \begin{cases} e^{l-1}_{n^i_m s}, & \text{if } r = n^j_m \text{ for some } m, \\ 0, & \text{otherwise,} \end{cases} \tag{4}$$

and

$$e^{l-1}_{rs} e^l_{ij} = (e^l_{ji} e^{l-1}_{sr})^* = \begin{cases} e^{l-1}_{r n^j_m}, & \text{if } s = n^i_m \text{ for some } m, \\ 0, & \text{otherwise.} \end{cases} \tag{5}$$

Hence, if $x \in K_l$ and $y \in K_{l-1}$ then $xy, yx \in K_{l-1}$. Suppose now that $x \in K_l$, $z \in A_{l-2}$. Then

$$xz = \lim_{m\to\infty} x(\sum_{i=1}^{m} e^{l-1}_{ii})z = \lim_{m\to\infty} (x \sum_{i=1}^{m} e^{l-1}_{ii})z.$$

We established that $x \sum_{i=1}^{m} e^{l-1}_{ii} \in K_{l-1}$ for every m hence $xz \in A_{l-2}$. Similarly, $zx \in A_{l-1}$ and we gather that $A_{l-1} = A_{l-2} + K_{l-1}$, $k \geq 2$, is an ideal in $A_l := A_{l-1} + K_l$ which is a C^*-subalgebra of $\mathcal{B}(H)$ by [6, 1.8.4]. We want now to prove $A_{l-1} \cap K_l = \{0\}$. Denote by B^m_l the finite dimensional C^*-algebra generated by $\{e^l_{ij} \mid 1 \leq i, j \leq m\}$. Then $K_l = \overline{\cup_{m=1}^{\infty} B^m_l}$. From

$$\| \sum_{i,j=1}^{m} \alpha_{ij} e^l_{ij} - \sum_{i,j=1}^{m} \alpha_{ij} e^l_{ij} \sum_{r=1}^{s} e^{l-1}_{rr} \| = \| \sum_{i,j=1}^{m} \alpha_{ij} e^l_{ij} \|$$

for every s we get $B^m_l \cap A_{l-1} = \{0\}$ for every m. Thus, the quotient map $A_l \to A_l/A_{l-1}$ is isometric on each B^m_l hence it is isometric on K_l and we conclude that $A_{l-1} \cap K_l = \{0\}$. A_l is a postliminal C^*-algebra since A_{l-1} and $A_l/A_{l-1} \sim K_i$ are postliminal C^*-algebras.

In this manner we construct inductively an increasing sequence $\{A_l\}_{l=0}^{\infty}$ of postliminal C^*-subalgebras of $\mathcal{B}(H)$ such that A_{l-1} is an ideal in A_l. It follows that $A := \overline{\cup_{l=0}^{\infty} A_l}$ is a postliminal C^*-subalgebra of $\mathcal{B}(H)$ whose greatest liminal ideal is A_0.

Set now $C_1 := K_1$, $C_2 := C_1 + K_2$. C_1 is an ideal in $\overline{C_2}$ hence C_2 is closed by [6, 1.8.4]. We have $C_1 \cap K_2 = \{0\}$ and $A_0 \cap C_2 = \{0\}$ since $(A_0 + K_1) \cap K_2 = \{0\}$ and $A_0 \cap K_1 = \{0\}$. We define inductively $C_l : C_{l-1} + K_l$. Then C_{l-1} is an ideal of the C^*-algebra C_l, $C_{l-1} \cap K_l = \{0\}$ and $A_0 \cap C_l = \{0\}$. From (3), (4) and (5) we find that $e_{ij}^l \to e_{ij}^{l-1}$, $1 \leq l \leq p$, $i, j \geq 1$ yields an isomorphism φ_p of C_p onto A_{p-1}. Obviously φ_{p+1} extends φ_p hence one gets an isomorphism φ from $C := \overline{\cup_{p=1}^{\infty} C_p}$ onto A that extends each φ_p. Let now $x \in A$, $x = \lim_{p\to\infty} x_p$ with $x_p \in A_p$, $p \geq 1$. Then $x_p = x_p^0 + x_p^p$ where $x_p^0 \in A_0$ and $x_p^p \in C_p$. From $\|x_p^0 - x_q^o\| \leq 2\|x_p - x_q\|$ we conclude that the Cauchy sequence $\{x_p^0\}_{p=1}^{\infty}$ converges to some $x^0 \in A_0$ hence $\{x_p^p\}_{p=1}^{\infty}$ converges to some $x^C \in C$ that satisfies $x = x^0 + x^C$. Now $x_p \to x_p^p$ is a homomorphism for each p therefore $x \to x^C$ is a homomorphism. We have $\|x^C\| \leq \|x\|$ and $\|x^0\| \leq 2\|x\|$. The quotient map $A \to A/A_0$ is isometric on each C_p hence it is isometric on C. It follows that $A_0 \cap C = \{0\}$ and the decomposition $x = x^0 + x^C$ is unique.

Now we can begin constructing the continuous field of C^*-algebras we need. Let $\{r_n\}$ be an enumeration of the set of rational numbers in $[0, 1]$. For an irrational number $t \in [0, 1]$ we define $A(t) := c_0(A)$ that is, the direct sum of A with itself $\aleph_0$ times. $A(r_n)$ is a C^*-subalgebra of $c_0(A)$ that is also a direct sum of copies of A except that at the n-th spot we insert C instead of A. Obviously all the fibers are mutually isomorphic postliminal C^*-algebras. The *-algebra Γ of the continuous vector fields consists of all the continuous functions $x : [0, 1] \to c_0(A)$ such that $x(t) \in A(t)$ for every $t \in [0, 1]$.

To show that $((A(t))_{t\in[0,1]}, \Gamma)$ so defined is a continuous field we must check that $\{x(t) \mid x \in \Gamma\} = A(t)$ for $t \in [0, 1]$. To this end let $t_0 \in [0, 1]$ and $\{a_n\} \in A(t_0)$. For $n \in \mathbb{N}$ let $f_n : [0, 1] \to [0, 1]$ be a continuous function such that $f_n(t_0) = 1$ and $f_n(r_n) = 0$ if $r_n \neq t_0$. Define $x(t) := \{f_n(t)a_n\}$, $t \in [0, 1]$. Then x is a continuous function from $[0, 1]$ to $c_0(A)$ such that $x(t) \in A(t)$ for $t \in [0, 1]$ i.e. $x \in \Gamma$. Moreover $x(t_0) = \{a_n\}$ and we have proved that we constructed a continuous field of C^*-algebras.

The greatest liminal ideal $B(t)$ of $A(t)$ is $c_0(A_0)$ when t is irrational. The greatest liminal ideal $B(r_n)$ of $A(r_n)$, $n \in \mathbb{N}$, is again a direct sum whose components are all equal to A_0 except the one at the n-th place that it is equal to K_1. Thus if

$x \in \Gamma$ satisfies $x(t) \in B(t)$ for every $t \in [0,1]$ then the n-th component of $x(r_n)$ must vanish since $A_0 \cap C = \{0\}$. It follows that for the restriction of our continuous field of C^*-algebras to any relatively open subset U of $[0,1]$ the family $(B(t))_{t\in U}$ together with $\{x \in \Gamma \mid x(t) \in B(t), t \in U\}$ does not form a continuous field of C^*-algebras.

References

1. R. J. Archbold and D. W. B. Somerset, *Transition probabilities and trace functions for C^*-algebras*, Math. Scand. **73** (1993), 81–111.
2. R. J. Archbold, D. W. B. Somerset and J. S. Spielberg, *Upper multiplicity and bounded trace ideals in C^*-algebras*, J. Funct. Anal. **146** (1997), 430–463.
3. E. Christensen and M.-D. Choi, *Completely order isomorphic and close C^*-algebras need not be *-isomorphic*, Bull. London Math. Soc. **15** (1983), 604–610.
4. E. Christensen, A. M. Sinclair, R. R. Smith, S. A. White and W. Winter, *Perturbations of nuclear C^*-algebras*, Acta Math. **208** (2012), 93–150.
5. J. Dixmier, *Sur les C^*-algèbres*, Bull. Soc. Math. France **88** (1960), 95–112.
6. J. Dixmier, *C^*-algebras*, North-Holland, Amsterdam, 1977.
7. J. M. G. Fell, *The structure of algebras of operator fields*, Acta Math. **106** (1961), 233–280.
8. P. Green, *Minimal primitive ideals of GCR C^*-algebras*, Proc. Amer. Math. Soc. **73** (1979), 209–210.
9. A. J. Lazar and D. C. Taylor, *Approximately finite dimensional C^*-algebras and Bratteli diagrams*, Trans. Amer. Math. Soc. **259** (1980), 599–619.
10. E. Michael, *Continuous selections. I*, Ann. of Math. **63** (1956), 361–382.
11. G. K. Pedersen, *C^*-algebras and their automorphism groups*, Academic Press, London, 1979.
12. J. Phillips, *Perturbations of C^*-algebras*, Indiana Univ. Math. J. **23** (1974), 1167–1176.
13. J. Phillips and I. Raeburn, *Perturbations of C^*-algebras, II*, Proc. London Math. Soc. **43** (1981), 46–72.

School of Mathematical Sciences, Tel Aviv University, Tel Aviv 69978, Israel
E-mail address: aldo@post.tau.ac.il